

Adult Neurology

Reed C. Perron, MD John T. Nasr, MD
Hugo N. Lijtmaer, MD Amrit K. Grewal, MD
Daniel R. Van Engel, MD Olga Noskin, MD
Kenneth A. Levin, MD Yamini Naidu, MD
Kenneth A. Citak, MD Daniel Berlin, MD, MSc
James T. Shammass, MD Fumin Tong, MD, PhD
Susan P. Molinari, MD Elena Zislin, PA-C

Pediatric Neurology

Peter L. Heilbronner, MD, PhD
Jennifer A. Cope, MD
Alexis M. Dallara-Marsh, MD
Mitchell Steinschneider, MD, PhD
Heather Weiner, APN

Neuro Rehabilitation Center

Kenneth A. Citak, MD
Medical Director

Managing Partner

Hugo N. Lijtmaer, MD

Chief Operations Officer

David T. Contento, FACMPE

John Jensen, PT

Director of Rehabilitation

SLEEP HYGIENE

- Sleep in a comfortable place (quiet, proper temperature, etc.)
- Keep a regular sleep-awake schedule. Get out of bed at the same time each morning.
- Avoid naps. Substitute light exercise for naps.
- Don't consume excessive alcohol or food close to bedtime. A light snack may help to induce sleep.
- Avoid caffeine, which may have a very prolonged stimulant effect.
- Put mental concerns "to sleep" by writing them down and setting them aside.
- Don't lie in bed unable to sleep for more than 30 minutes. Get out of bed, do something relaxing, and return to bed only when sleepy.
- Don't worry about the impact of missing sleep. In most cases, the mind and body will eventually get the rest that they need, and there seems to be little harm in the waiting.